



One-Pass/Two-Pass – South

DuPont™ Canopy® herbicide delivers:

- Residual activity to help maximize early-season control of tough weeds, allowing your postemergence application to be made closer to canopy
- Two modes of action to help prevent and manage weed resistance, such as glyphosate resistant Palmer pigweed and marestail

Application Information

- For Enlist E3™ soybeans, follow with an in-crop application of Enlist Duo® herbicide, Enlist One® herbicide, Liberty® herbicide, or Enlist One + Liberty
- For glyphosate-tolerant soybeans, follow with an in-crop application of DuPont™ Abundit® Edge herbicide (glyphosate)
- For glufosinate-tolerant soybeans follow with an in-crop application of glufosinate
- As needed, use products such as DuPont™ Synchrony® XP herbicide following the in-crop application to broaden weed-control spectrum
- In the Southern states, do not apply more than 3.5 ounces to soils with an aggregate pH greater than 7.0
- Refer to the label for pH restrictions in your state
- DO NOT apply Canopy postemerge to emerged soybeans

Weeds controlled – preemergence

| | | |
|----------------------------------|-----------------------------------|----------------------------------|
| S Barnyardgrass | C Lambsquarters | C Ragweed, giant |
| C Beggarweed, Florida | C Marestail (horseweed) | C Sicklepod |
| S Burcucumber | C Morningglory spp. | C Sida, prickly (teaweed) |
| C Cocklebur | C Mustard, wild | S Signalgrass, broadleaf |
| C Copperleaf, hophornbeam | S Nutsedge: purple, yellow | C Shepherd's purse |
| S Crabgrass | S Panicum: fall, Texas | C Smartweed, Pennsylvania |
| S Foxtail, species | C Pigweed spp. | C Spurge, spotted |
| C Henbit | C Poinsettia, wild | C Velvetleaf |
| S Johnsongrass, seedling | C Ragweed, common | C Yellow rocket |

C – Controlled | **S** – Suppressed

See Label for full details

| Canopy 2-Pass (PRE by POST) use rates | Preemergence Broadcast use rates (oz/A) | |
|--|---|------------------|
| | 0.5 – 4% organic matter in soil | |
| Soil texture | Southern States* | Northern States* |
| Any** | 3 – 3.5 | 2.25 |
| Coarse: Loamy sand, sandy loam | 4 – 6 | 3 – 5 |
| Medium: Loam, silt loam, silt, sandy clay loam | 4 – 6 | 3 – 6 |
| Fine: Silty clay loam, clay loam, clay | 6 – 8 | 5 – 7 |

Canopy may need to be followed by a post application of Synchrony XP, Abundit Edge, Durango® DMA® herbicide, or other post products allowed by the trait package

* Refer to the Canopy label for a complete list of Northern and Southern states

** 3.5 ounces per acre is the maximum rate that may be used on soils with a composite pH greater than 7.0, no chlorimuron-ethyl containing herbicide such as Synchrony® XP should be used in sequential treatments

| Crop | Rotation (Months) | | Crop | Rotation (Months) | |
|--------------|-------------------|----------------|----------------------|-------------------|----------------|
| | A ^{1,2} | B ³ | | A ^{1,2} | B ³ |
| Soybeans | Anytime | Anytime | Cotton | 10 | 10 |
| Wheat | 4 | 4 | Peanuts | 8 | 18 |
| Clover | 12 | 12 | Tobacco (transplant) | 10 | 10 |
| Corn, Field* | 9/10** | 10 | Other | See Label | |

¹ States of AL, AR, FL, GA, LA, MS, TX;

² States of KY, MO bootheel, NC, OK, SC, TN, TX

– Use rate less than 10.0 oz/A; If soil above 7.0 pH and Canopy rate is greater than 3.5 oz/A, see label for additional restrictions

³ VA and WVA

* Field Corn is defined as, grain, silage, popcorn, see label for seed corn

** Rotation is 9 mo if the Canopy rate does not exceed 6 oz

Burndown – South

DuPont™ Canopy® herbicide delivers:

- Start clean with Burndown and residual activity of Canopy to keep fields clean up to planting
- Two modes of action to help prevent and manage weed resistance, such as glyphosate resistance marestail
- Reliable burndown of winter annuals and early-season weeds, even under cool conditions, for a wide application window

Application Information

- Canopy may be applied in fall, anytime post harvest, up to at-plant.
- For residual control till planting, Canopy may be applied anytime prior to planting up to 45 days before

Weeds controlled – burndown¹

| | |
|---|---|
| C Annual grasses | C Mustard, tansy, wild |
| C Buttercup, small flower | C Pennycress, field |
| C Deadnettle, purple | C Shepherds purse |
| C Garlic, wild ² | C Speedwell, field, purslane |
| C Henbit | C Thistle, Canada (above ground portion) |
| C Lettuce, prickly | C Yellow rocket |
| C Marestail (horseweed) ² | |

¹ Canopy used for burndown must include either a crop oil concentrate (COC) or NIS.

² The addition of 8 oz/A of 2,4 D LVE is required for all Canopy rates


C – Controlled | **S** – Suppressed

See Label for full details

| Canopy Burndown (after harvest up to planting) | Burndown with residual broadcast rate (oz/A) – Soil pH of 7 or less* | |
|---|---|------------------|
| | 1.5 – 4% organic matter in soil | |
| Soil texture | Southern States** | Central States** |
| Coarse: Loamy sand, sandy loam | 3.5 – 5 | 3 – 5 |
| Medium: Loam, silt loam, silt, sandy clay loam | 3.5 – 6 | 3 – 6 |
| Fine: Silty clay loam, clay loam, clay | 5 – 7 | 5 – 7 |

* No soil pH restriction for Central States – 2.25 oz/acre | No soil pH restriction for Southern States – 2.25 – 3.5 oz/acre

** Refer to the Canopy label for a complete list of Northern and Southern states



Visit us at corteva.us

®/™ Trademark of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use, and product warranty and liability limitations, which must be read before using the product(s). Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label direction and precautions for use when using any pesticide alone or in tank-mix combinations.

Not all products are registered for sale or use in all states. Contact your state pesticide regulatory agency to determine if a product is registered for sale or use in your state.

Enlist E3™ soybeans were jointly developed by Dow AgroSciences and MS Technologies. Enlist One® and Enlist Duo® herbicides is the only 2,4-D product authorized for use in Enlist crops. Consult Enlist herbicide labels for weed species controlled. Always read and follow label directions. Liberty and LibertyLink are registered trademarks of BASF. ©2020 Corteva. CA01-137-068 COR (04/20)