DuraCor® Herbicide Handling Recommendations

DuraCor® herbicide is a suspension concentrate (SC) formulation that allows for a low and flexible use rate and the ability to mix well with fertilizer. As with all SC formulation types, it is recommended to shake or mix the product prior to application for best results. Below are best management practices for mixing DuraCor herbicide.

	4x1 Gallon	Drum (30 Gallon)	Tote (250 Gallon)	Bulk
BEGINNING of Application Season (BEFORE first application from product container after opening for use)	Shake Well Before Use	Mix for 30 Minutes	Mix for 30 minutes	Recirculate the tank (3 volume turns)
DURING Application Season	Shake Well Before Each Use	Mix for 30 Minutes Per Month	Mix for 30 minutes Per Month	Recirculate the tank (3 volume turns) Per Month
AFTER Application Season	No Mixing Required as Product Will Freeze ~25°F.			

Talk to your Corteva Agriscience™ Range & Pasture Specialist for additional details.

Duracorherbicide.com







Visit us at rangeandpasture.com

Under normal field conditions this product is non-volatile. DuraCore has no grazing or haying restrictions for any class of livestock, including lactating dairy, cows, horses (including lactating mares) and meat animals prior to slaughter. Label precautions apply to forage treated with DuraCor to manure and urine from animals that have consumed treated forage entractemarks of Dow AgroSciences, DuPort or Ploneer and their affiliated companies or respective owners. DuraCor is not registered for sale or use in all states. Contact your state pesticide regulatory agency to determine if a product is registered for sale or use in your state pesticide regulatory agency to determine if a product is registered for sale or use in your state pesticide regulatory agency to determine if a product is registered for sale or use in your state. Always read and follow label directions. © Corteva CR45 640 022 (06/20) 010-58917