

Rezuvant[®]

Arylex[™] active

HERBICIDE

Cross-spectrum weed control solution for wheat and barley

Rezuvant[®] herbicide with Arylex[®] active combines the next generation of broadleaf weed control with an industry-leading graminicide in one convenient premix, providing cereal growers with a straightforward solution for conquering tough weeds.

Benefits

- Cross-spectrum control of broadleaf and grass weeds in one, straightforward solution
- Superior performance against troublesome grass weeds such as Italian ryegrass, foxtail, Persian darnel and wild oats, including Group 2-resistant biotypes
- An elevated load of fluroxypyr plus the power of Arylex[®] active delivers consistent control of hard-to-manage broadleaf weeds like kochia, wild buckwheat, giant ragweed and common lambsquarters

Features

- 3 sites of action – pinoxaden (Group 1), fluroxypyr (Group 4) and Arylex (Group 4)
- Application flexibility from 3-leaf to flag leaf emergence
- Rainfast in 1 hour
- Stable across a wide range of temperatures, including for the early-season applications in the northern U.S. grain production areas
- Crop rotation the following spring to all major crops, except for lentils
- Convenient tank mix capabilities with other herbicides (see label for guidelines)
- Flexible for performance in all situations

Application recommendations

Labeled application rate: 16.4 fl oz/A

- Apply Rezuvant to spring wheat, winter wheat and barley from 2-leaf to flag leaf emergence
- For best results, apply Rezuvant when susceptible broadleaf weed seedlings are actively growing and less than 4 inches tall
- Do not harvest treated crop for grain within 60 days after application
- Do not apply closer than 30 days before cutting of hay
- Livestock grazing permitted on treated crops 30 days following application

Crop	Rotational interval (months) ¹
Barley, wheat (including spring, winter, and durum)	0
Canola, cotton, millet, popcorn, rice, rye, seed corn, soybean, sugarcane, sunflower, field corn, oats, sorghum, sweet corn	4
Alfalfa, brassica (cole) leafy vegetables, camelina, chickpea, clover, dry bean, flax, mustard, peanut, peas (dry and succulent), safflower, sugar beet	9
Potato ²	10
Other crops not listed	15

1. Minimum number of months that must pass before planting other crops after application of Rezuvant herbicide.

2. For rotation to potatoes, precipitation (including irrigation) must be greater than 16.0 inches during the 10 months following application of Rezuvant. Otherwise, rotation to potatoes is recommended 15 months following application.

Grass weeds controlled

Barnyardgrass	Oats, volunteer
Canarygrass	Oats, wild
Foxtail, giant	Persian Darnel
Foxtail, green	Ryegrass, Italian
Foxtail, yellow	Windgrass
Millet, wild proso	

Broadleaf weeds controlled

Alfalfa seedlings, volunteer	Morningglory
Catchweed bedstraw (cleaver)	Nightshade, eastern black
Chickweed, common	Nightshade, hairy
Cocklebur	Nightshade, cutleaf
Cressleaf groundsel	Pigweed, redroot
Flax, volunteer	Plantain, common
Flixweed	Prickly lettuce
Fumitory	Puncturevine
Hairy vetch	Purple deadnettle
Hempnettle, common	Purslane, common
Henbit	Ragweed, common
Horseweed (marestail)	Ragweed, giant
Kochia	Sunflower, common
Lambsquarters, common	Velvetleaf
Mallow, common	White clover
Mallow, Venice	Wild buckwheat

Visit us at corteva.us



This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use, and product warranty and liability limitations, which must be read before using the product(s). Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label direction and precautions for use when using any pesticide alone or in tank-mix combinations.

Rezuvant[®] is not registered for sale or use in all states. Contact your state pesticide regulatory agency to determine if a product is registered for sale or use in your state. Always read and follow label directions.

©2024 Corteva. COR (04/24)