



HERBICIDE

HERBICIDE

GREAT POSTEMERGENCE TANK MIX PARTNERS FOR MORE CONSISTENT CONTROL OF TOUGH BROADLEAF WEEDS IN GLYPHOSATE TOLERANT SOYBEANS

Benefits of Multiple Modes of Action

- Helps combat increased glyphosate tolerance
- Improves speed of knockdown on key broadleaf weeds
- Provides more consistent control of larger broadleaf weeds than glyphosate alone

Benefits of Residual Chemistry

- Reduces the likelihood for a third application
- Minimizes late season flushes of broadleaf weeds
- Allows for better timing of subsequent glyphosate applications

Application Timing

- For best control apply when weeds are no larger than 4-6" tall
- Durango[®] DMA[®] may be post-applied from emergence through flowering (R2 soybean stage)
- FirstRate[®] may be applied any time prior to the R2 (full flower) growth stage of soybeans
- Offers excellent crop safety throughout application window

Control Target Weeds with FirstRate[®] and Durango[®] DMA[®]



Ragweed, common and giant



Morningglory



Velvetleaf



Yellow Nutsedge




Marestail



Cocklebur

FirstRate[®] and Durango[®] DMA[®] – Complementary Modes of Action



Tank Mix Recommendation

Durango DMA ¹ or Duramax ^{®*}	24 - 48 oz/A
FirstRate ²	0.3 - 0.4 oz/A
AMS	2 lb/A
NIS	1 - 2pt/100 gal. (May be added to improve control of large weeds or weeds under drought stress)

¹Total in-crop applications not to exceed 72 oz, 48 oz, in a single application.

²Total in-crop application not to exceed 0.6 oz, either as a single application or as a total of sequential of post emergence applications. The maximum cumulative application rate from preplant, preemergence, and/or postemergence must not exceed 1.05 oz per growing season.

For more information about Durango[®] DMA[®] or FirstRate[®] herbicide contact your local Dow AgroSciences sales representative or visit www.dowagro.com/usag.