



DESIGNED FOR USE IN ROUNDUP READY® CORN

1st Pass - Soil Applied

SureStart® Herbicide Use Rates

| Soil Texture | Soil OM < 3% | Soil OM ≥ 3% |
|--------------|----------------|-----------------|
| Coarse | 1.5 - 2.0 pt/A | 1.5 - 2.0 pt/A |
| Medium | 1.5 - 2.5 pt/A | 1.75 - 3.0 pt/A |
| Fine | 2.0 - 3.0 pt/A | 2.0 - 3.0 pt/A |

- Apply early preplant up to 30 days before planting, preplant incorporated, preemergence or to spike stage corn. For maximum residual, apply as close to planting as possible
- Effective in all tillage systems
- Three non-glyphosate modes of action for improved resistance management
- Provides 5-6 weeks of residual control, designed to be followed by a post application of glyphosate when weeds reach 4-6" in height
- Do not use with Counter insecticide; plant corn at least 1.5" deep
- Doesn't contain atrazine – may be used in atrazine restricted areas

2nd Pass - Post Applied

- Apply Durango® DMA® herbicide at 32 oz/A to 4-6" weeds
- Optimal weed control timing for maximum yield protection

Early Post Tankmix

- SureStart® herbicide + Durango® DMA® tank mixture may be applied from spike to 11" tall corn
- Apply to weeds 1-2" in height
- Do not use liquid fertilizer as the carrier

| | |
|---------------|-------------------|
| SureStart® | 2.0 – 2.5 pt/acre |
| Durango® DMA® | 32 oz/acre |
| AMS | 2.5 lbs/acre |

WEEDS CONTROLLED:

- barnyardgrass
- Canada thistle¹
- cocklebur
- crabgrass
- fall panicum
- field sandbur
- foxtail species
- jimsonweed
- Johnsongrass, seedling
- kochia
- lambsquarters
- marehail
- morningglory, annual
- nightshade species
- pigweed species
- prickly sida
- ragweed, common
- ragweed, giant
- shattercane
- smartweed
- sunflower
- velvetleaf
- Venice mallow
- waterhemp
- witchgrass
- woolly cupgrass
- yellow nutsedge

¹Burndown activity in minimum and no-till corn only
Partial list of weeds controlled, always refer to label