



# PerfectMatch®

**POWER TO CONTROL 86 BROADLEAF WEEDS AND 13 GRASS SPECIES WITH JUST ONE HERBICIDE PRODUCT IN YOUR TANK**

## HERBICIDE

PerfectMatch® herbicide combines the proven broadleaf weed control of WideMatch® herbicide with the leading grass control in spring wheat (including durum) and winter wheat to deliver the broadest control in one new product.

### NOTEWORTHY FEATURES

- Two modes of action in three active ingredients help manage herbicide resistance in wheat, including durum, while controlling larger, hard-to-control weeds.
- PerfectMatch is the simplest solution for your customers to control the broadest spectrum of broadleaf weeds and grasses.
- This herbicide is a tank-mix-compatible Suspo - Emulsion (SE) formulation, so you can combine your favorite approved adjuvant or partners.



Active Ingredient	Mode of Action	Site of Action
Clopyralid	Growth regulator	Group 4
Fluroxypyr	Growth regulator	Group 4
Pyroxsulam	ALS inhibitor	Group 2

### APPLICATION RECOMMENDATIONS


- Application rate: 10 GPA carrier or greater.
- The best results are obtained when grass weeds are treated at the two-leaf to the two-tiller stage of growth and before broadleaf weeds are more than 2 inches tall or 2 inches in diameter.
- Apply with a high-quality nonionic surfactant at a rate of 0.25 percent to 0.5 percent v/v (1 to 2 quarts per 100 gallons of spray solution).
- Ammonium sulfate (AMS) is recommended, as well, or urea ammonium nitrate (UAN) fertilizer.
- When PerfectMatch is tank-mixed with emulsifiable concentrate (EC) formulations above 6 fl.oz. /A, do not use a surfactant.
- Applications of PerfectMatch are rainfast within four hours after application.

Crop <sup>1</sup>	Rotation Interval
Wheat, triticale	1 month
Barley, camelina, canola (rapeseed), cotton, field corn, flax, grasses, millet, mustard, oats, popcorn, seed corn, sugarbeet, sweet corn	9 months
Alfalfa, dry beans, peas (dry and succulent <sup>2</sup> ), grain sorghum, onions, safflower, soybeans, strawberries, sunflower	10.5 months
Chickpeas, cole crops (brassica species), lentils, potatoes (including potatoes grown for seed), and broadleaf crops grown for seed	18 months
Other crops not listed	18 months

1. A field bioassay is recommended prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 18 months following application. Crop rotation intervals for all states except Idaho, Nevada, Oregon, Utah, and Washington.
2. For rotation to field peas in 10.5 months, precipitation must be greater than 7.0 inches during the 10.5 months following application of PerfectMatch™ herbicide and greater than 5.5 inches during the June 1 through August 31 time period following application. Otherwise, rotation to field peas is recommended 18 months following application.

## CANADA THISTLE CONTROL

6 to 8 Weeks After Treatment



1. One- to four-leaf stage of growth.
2. For best control, apply up to stage of growth.
3. Including herbicide-tolerant canola varieties except CLEARFIELD® (imidazolinone-tolerant) canola.
4. Control may be reduced when application is made after bolting.
5. Less than 2 inches tall. For control of lambsquarters 2 inches tall, 0.25 2,4-D ester or MCPA ester. For control of Russian thistle 2 inches tall, with 0.25 2,4-D ester.
6. For best control or suppression, apply at the stage of growth. PerfectMatch™ herbicide will control the initial top growth and inhibit regrowth during the season of application (season-long control).
7. For best control or suppression, apply from rosette to bud (pre-flower) stage of growth.
8. Includes resistant biotypes. Best control is achieved when weeds are at least 1 inch tall.



For more information about Dow AgroSciences or any of our crop protection products for small grains, visit [dowagro.com/usag](http://dowagro.com/usag).

GRASS WEEDS	CONTROLLED (C) OR SUPPRESSED (S)
Barley, foxtail	S
Barnyardgrass	C
Brome, downy	S
Brome, Japanese	C
Cheat	C
Darnel, Persian	C <sup>1</sup>
Foxtail, green	S
Foxtail, yellow	C <sup>1</sup>
Oats, wild	C
Quackgrass	S
Ryegrass, Italian	C

BROADLEAF WEEDS	CONTROLLED (C) OR SUPPRESSED (S)
Bindweed, field	S
Buckwheat, wild	C
Canola, volunteer (wild turnip)	C <sup>3</sup>
Chamomile, false (scentless)	C
Cocklebur, common	C
Daisy, oxeye	S
Dandelion	C
Flax, volunteer	C
Hawksbeard, narrowleaf	C
Horsetail, field	S
Horseweed	C
Kochia	C <sup>8</sup>
Lambsquarters, common	C <sup>5</sup>
Mallow, common	S
Mustard, wild	C
Nightshade species	C <sup>7</sup>
Pigweed, redroot	C
Ragweed, common	C <sup>2</sup>
Ragweed, giant	C <sup>2</sup>
Sowthistle, annual	C
Sowthistle, perennial	S <sup>6</sup>
Starthistle, yellow	C
Sunflower	C
Thistle, Canada	C <sup>6</sup>
Thistle, Russian	C <sup>5</sup>
Wallflower, bushy	C <sup>4</sup>
Wormwood, biennial	C



Dow AgroSciences

Solutions for the Growing World