Intrepid Edge®

INSECTICIDE

Effective Control of Citrus Leafminers and Asian Citrus Psyllid


Features and Benefits

- · Two Unique modes of action
 - » Methoxyfenozide (Group 18)
 - » Spinetoram (Group 5)
- Easy to mix and use Suspended Concentrate (SC) formulation
- · Fast knockdown, death in hours
- Effective control of key citrus pests

Restrictions Citrus

- · Do not apply within 1 day of harvest
- Do not apply more than
 12 fl oz/A per application
- Do not apply more than 48 fl oz/A per year
- · Minimum treatment interval is 7 days apart
- Do not apply more than 3 applications per year
- Do not apply to citrus nurseries or citrus in greenhouses

Intrepid Edge® insecticide will provide excellent control of citrus leafminers Chart 1. Citrus leafminer control:



Control all life stages of Asian Citrus Psyllid

Chart 2. Asian citrus psyllid egg control




Chart 3. Asian citrus psyllid nymph control





Chart 4. Asian citrus psyllid adult control



All Treatments Contain a 0.25% MSO/OS John Curtis, 2016 Port St. Lucie, FL

Application Rates Citrus

Pest	Intrepid Edge® Rate (fl oz/A)
Leafrollers	
Citrus Leafminer	
Orange Dog	
Peelminer	
Cutworms	6-12
Light Brown apple moth	, . <u></u>
Orange Tortix	
Western	
Tussock Moth	
Asian Citrus Psyllid*	6-8

Visit us at corteva.us

pesticide regulatory agency to determine if a product is registered for sale or use in your state. Always read and follow label directions. ©2019 Corteva CA01-911-021 COR (06/19)

