

# Count on consistent control of grasses and broadleaf weeds in corn

Keystone® NXT herbicide is the dependable soil-applied solution for corn growers seeking consistent control of a wide spectrum of grasses and small-seeded broadleaf weeds. including foxtails, barnyardgrass, crabgrass, waterhemp, lambsquarters, pigweeds, nightshade and many more.

### Providing dependable and flexible control

- Keystone NXT offers one-pass control of many tough annual grasses and broadleaf weeds.
- Dual modes of action provide control of herbicide-resistant weeds supporting glyphosate stewardship programs.
- Keystone NXT requires only 1/4 inch of rain for activation.
- It can be applied from early preplant to early postemergence.
- It fits conventional, reduced-till and no-till corn production systems.

## **Target application timing**







Preplant Preemergence

Postemergence

Up to 11 inch corn

## Use rates (qt./A) by soil texture and organic matter content in conventional tillage systems

Soil	Broadcast rate per Acre (qt.)* Soil Organic Matter Content	
texture	Less than 3%	More than 3%
Coarse	1.4 qt.	1.7 qt.
Medium	1.7 – 2.4 qt.	2.3 - 2.6 <sup>†</sup> qt.
Fine	2.3 - 2.6 <sup>†</sup> qt.	2.3 - 3.0 <sup>†</sup> qt.

<sup>\*</sup>Use the higher rate in the rate range in areas of heavy weed infestation.

For more information about Keystone NXT herbicide, visit KeystoneNXT.com, call 800-258-3033 or contact your local Corteva Agriscience territory manager.

#### Receive a broad spectrum of control

Keystone NXT combines the power of acetochlor and atrazine for consistent control of annual grasses and small-seeded broadleaf weeds. See product label for application rates and instructions.

Weeds controlled or partially controlled			
Grasses & Sedges			
Barnyardgrass	Nutsedge, yellow <sup>1,2</sup>		
Crabgrass spp.	Oats, wild		
Crowfootgrass	Panicum, browntop		
Cupgrass, southwestern	Panicum, fall		
Cupgrass, woolly*	Panicum, Texas <sup>3*</sup>		
Foxtail, giant	Rice, red		
Foxtail, green	Sandbur, field*		
Foxtail, robust (purple, white)	Shattercane*		
Foxtail, yellow	Signalgrass, broadleaf <sup>3</sup>		
Goosegrass	Sprangletop, red		
Johnsongrass, seedling*	Wheat, volunteer		
Millet, foxtail	Witchgrass		
Millet, wild proso*			

Broadleaves		
Beggarweed, Florida	Nightshade, hairy	
Carpetweed	Pigweed, spp.	
Cocklebur <sup>2</sup>	Purslane, common	
Galinsoga	Pusley, Florida	
Groundcherry, annual	Ragweed, common	
Groundcherry, cutleaf	Ragweed, giant*	
Henbit	Sicklepod*	
Jimsonweed	Sida, prickly	
Kochia	Smartweed spp.	
Lambsquarters, common	Sunflower, common*	
Morningglory spp. <sup>2</sup>	Velvetleaf <sup>2</sup>	
Mustard spp.	Waterhemp, tall	

Nightshade, black

Visit us at corteva.us





<sup>&</sup>lt;sup>†</sup>On highly erodible soils with less than 30% plant residue, do not apply more than 2.5 quarts of Keystone® NXT herbicide per acre in a single preemergence application.

<sup>\*</sup> Partial Control

<sup>&</sup>lt;sup>1</sup> Preplant incorporate for improved control.

<sup>&</sup>lt;sup>2</sup> Use the higher rate in the specified application rate range. Control of these weeds can be erratic, especially under dry conditions. Additional atrazine and/or sequential herbicide applications may be needed for complete control.

<sup>&</sup>lt;sup>3</sup> Best control is achieved when Keystone NXT is applied within 5 days of planting and rainfall occurs shortly after application or mechanical incorporation is used to activate the herbicide. If rainfall does not occur within 7 days after application, shallow cultivation will enhance activity. Excessive rainfall after application may reduce control. Under adverse weather conditions and/or heavy infestations, a cultivation or follow-up herbicide application may be needed.