

The Advantage of Transline® Herbicide

Transline®
HERBICIDE

The Advantage of Transline® Herbicide

Control broadleaf weeds and woody brush species in non-crop areas, forest sites, industrial manufacturing and storage sites, rights-of-way, and wildlife openings, including grazed areas on these sites, tree plantations, and rangeland and permanent grass pastures.

Transline has excellent selectivity, allowing it to be used over many conifers and hardwood species. It is safe on warm- and cool-season right-of-way grasses. Transline offers excellent residual control and is rainfast within two hours. It offers tank-mix compatibility and has low volatility.

Benefits

- Excellent control of many tough western noxious invasive weeds, including knapweeds and thistles, including infestations in and around many pine trees. Unlike most other herbicides, including Redeem® Range & Pasture, Transline® herbicide can be applied over the top of pine trees.
- Use 2/3 - 11/3 pints/acre of Transline for selective control of thistle and knapweeds around trees. A non-ionic surfactant containing at least 80% active ingredient at 1/4 -1/2 percent can be added to improve herbicide wetting and penetration.
- Transline is available in 1/2 gallon containers and 2-1/2 gallon containers as well as the Continuum™ prescription control and container management system. One-half gallon containers are very popular for small end users that used to use Redeem Range & Pasture herbicide.

Control of High Anxiety Invasives

Treatment Method Timing

Spotted Knapweed

Apply herbicides when plants are at the rosette to the mid-bolting stage or fall regrowth.

Russian Knapweed

Apply herbicides, from bud to mid-flower growth stages or on dormant plants in the fall. Treatments applied in the fall will be made over plants where the foliage is dead and brown. Apply when the plants are actively growing. When treating bi-annual thistles, it is best to treat them while they are in the rosette stage in the fall or in the spring before plants flower. Canada thistle is best controlled during fall regrowth or during bolting to early bud stage.



Visit us at vegetationmgmt.com

®/™Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners. When treating areas in rangeland, wildlands, and around roadside or utility rights-of-way that are or will be grazed, hayed or planted to forage, important label precautions apply regarding harvesting hay from treated sites, using manure from animals grazing on treated areas or rotating the treated area to sensitive crops. Consult the label before purchase or use for full details. State restrictions on the sale and use of Transline apply. Consult the label before purchase or use for full details. Always read and follow label directions. ©2019 Corteva Agriscience CV01 113 020 (05/19) 010-51149



Broadleaf Weeds Controlled

Acacias	Hawkweed, Orange	Ragweed, Common
Artichoke, Jerusalem	Hawkweed, Yellow	Ragweed, Giant
Buckwheat, Wild	Horseweed	Salsify, Meadow (Goatsbeard)
Buffalobur ¹	Jimsonweed	Sicklepod
Burdock, Common	Knapweed, Difuse	Smartweed, Green ¹
Chamomile, False (Scentless)	Knapweed, Russian ¹	Sorrel, Red
Chamomile, Mayweed (Dogfennel)	Knapweed, Spotted	Sowthistle, Annual
Clover, Black Medic	Ladysthumb ¹	Sowthistle, Perennial ¹
Clover, Hop	Lettuce, Prickly	Starthistle, Yellow
Clover, Red	Locoweed, White	Sunflower (Common And Wild)
Clover, White	Locoweed, Lambert	Teasel, Common
Cocklebur, Common	Marshead	Thistle, Artichoke
Coffeeweed	Mesquite	Thistle, Bull
Cornflower (Bachelor Button)	Nightshade, Eastern Black	Thistle, Canada (Rosette To Bud)
Dandelion	Nightshade, Cutleaf	Thistle, Musk (Rosette To Bud)
Dock, Curly	Nightshade, Hairy	Thistle, Italian
Groundsel, Common	Oxeye, Daisy	Vetch
Hawksbeard, Narrowleaf	Pineappleweed	



Woody Plants and Vines Controlled

Eastern Redbud	Knapweed Diffuse	Thistle, Artichoke
Kudzu	Knapweed Russian ¹	Thistle, Canada (Rosette to Bud)
Locust (Spp)	Knapweed Spotted	Thistle, Italian
Mimosa (Silktree)	Starthistle, Yellow	Thistle, Musk (Rosette to Bud)
Wisteria		

