



# COMBINE PASTURE WEED CONTROL WITH DRY FERTILIZER

## FERTILIZE YOUR GRASS, NOT YOUR WEEDS – CONVENIENTLY.

### Benefits:

- Save time and extra application cost — it's one trip
- Fewer weather delays
- Reduced risk of drift
- No spray equipment needed
- Self-apply on your schedule if custom application is not available
- Fits large or small operations

### How it works:

Impregnation is the process of applying a concentrated herbicide solution on dry fertilizer granules during the blending process. To get an adequate rate properly distributed, apply GrazonNext® HL or Chaparral™ specialty herbicides with at least 200 pounds of dry fertilizer per acre.

Spreader trucks or fertilizer buggies apply the dry product as they would normally apply dry fertilizer alone. From there, rainfall puts the herbicide-fertilizer solution into the soil. GrazonNext HL and Chaparral are soil active herbicides absorbed by weeds through the roots. Weed control is almost entirely dependent on the soil residual activity of the herbicide and root uptake by the weeds.

Expectations for weed control should be less when compared to foliar applications. However, research trials have demonstrated up to 85% control while many ranchers have reported better results.



## States with 2(ee) Recommendation for GrazonNext® HL or Chaparral™ impregnation on dry fertilizer:



### Tips for application:

- Apply GrazonNext® HL at a rate of 2 – 2.1 pt/A or Chaparral™ at a rate of 2 – 3.3 oz/A
- Include a dye to alert all users to the presence of the herbicide
- Apply at least 200 pounds of fertilizer per acre
- Apply when weed species are young, 4-6 inches tall in most cases. Older more mature weeds may be harder to control.
- Do not use Chaparral on bahiagrass if damage to bahia cannot be tolerated
- The mixing and application equipment must be dedicated for use only on grasslands

### Follow good stewardship:

GrazonNext HL and Chaparral have no grazing or haying restrictions for any class of livestock, including lactating dairy cows, horses (including lactating mares) and meat animals prior to slaughter. However, label precautions do apply to forage treated with GrazonNext HL and Chaparral and to manure from animals that have consumed treated forage within the last three days. Consult the label for details.