

# Dimension<sup>®</sup>

## SPECIALTY HERBICIDE


The leading turf and ornamental herbicide that provides preemergence and early postemergence control of crabgrass and other important grassy and broadleaf weeds.

### Effective control of crabgrass

Dimension<sup>®</sup> specialty herbicide makes controlling crabgrass easier because it is the only preemergence herbicide that offers excellent postemergence control. It also is proven effective for more than 45 other grassy and small-seeded broadleaf weeds. Eliminating weeds before they emerge means fewer customer complaints.

### Long application window

By providing both pre- and early postemergence control, Dimension<sup>®</sup> specialty herbicide offers a longer application window than other preemergence herbicides, which allows more time to make applications. Applicators can confidently use Dimension from early to midseason without having to worry about switching products after crabgrass has germinated.



### Application flexibility and nonstaining formulations

Dimension<sup>®</sup> specialty herbicide can be applied to turf, landscapes and bare ground – making it the ideal wall-to-wall solution. And no matter which formulation of Dimension is applied (liquid, WSP, granular and on-fertilizer), there are no concerns about it staining your equipment or your customers' structures and driveways yellow.

### KEY FACTS

- Effective control of crabgrass and more than 45 other grassy and broadleaf weeds
- Reduces hand-weeding and lowers labor costs in landscape and nursery settings
- Excellent preemergence and early postemergence crabgrass control
- Available in multiple formulations: liquid, WSP, granular and on-fertilizer
- Labeled for use on:
  - Established cool- and warm-season turfgrass
  - More than 440 ornamental plants
  - Over-the-top application on 142 field-grown ornamentals and 125 container-grown ornamentals

### Exceptional plant safety

Weed control only matters if turf and ornamentals continue looking vibrant after application. Dimension<sup>®</sup> specialty herbicide has proven exceptional turf and ornamental safety on both established cool- and warm-season turfgrasses as well as more than 440 ornamental plants. The broad label enables more confidence that the weeds, not desirable plants, are controlled.

### Target weeds

- Bittercress
- Chickweed
- Crabgrass
- Goosegrass
- Oxalis
- *Poa annua*
- Spurge



### When to use Dimension® specialty herbicide

Dimension® specialty herbicide is the leading turf and ornamental herbicide providing pre- and early postemergence control of crabgrass as well as other important grassy and broadleaf weeds. It can be applied before weeds emerge and early postemergence, up to the five-leaf stage for crabgrass, when applied in areas where warm-season turfgrasses are predominant.

Dimension can be tank-mixed with Defendor® specialty herbicide and Gallery® specialty herbicide or other herbicides for an even broader spectrum of weed control. Please refer to the label for information on the other products compatible for tank-mixing with Dimension.

### How to use Dimension® specialty herbicide

Choose the formulation that works best for your operation – liquid, granular, wettable powder or on-fertilizer – and use it according to label directions for your specific region of the country and particular program conditions.

When using a liquid formulation, apply with a sufficient volume of water carrier to provide a uniform spray distribution. Be sure that treated areas receive a ½ inch of rainfall or irrigation after treatment. Granular treatments do not require immediate watering; however, best results will be obtained if irrigation or rainfall occurs within seven days of the application.

### Application rate

For preemergence grass and broadleaf weed control, apply Dimension® 2EW specialty herbicide as a single or sequential application at 1 to 2 pints (0.25 to 0.5 lb. active ingredient) per acre. Applicators may choose to make a single application or sequential applications of 1 to 2 pints per acre at five- to 10-week intervals based on various factors listed on the label.

For postemergence control, when applied at 2 pints (0.5 lb. active ingredient) per acre, Dimension 2EW has demonstrated crabgrass control through the three- to five-tiller stage of growth in the western, southern and transition regions, where warm-season turfgrasses are the predominant species.

In regions where cool-season turfgrasses are the predominant species, early postemergence crabgrass control is obtained when Dimension 2EW is applied before the initial tiller stage of crabgrass (less than five leaves per plant), which generally corresponds to the time when crabgrass seedlings are easily observed in lawn or turf.

