

HERBICIDE



Stewardship guidelines

NativeKlean™ herbicide is labeled for use **only** in native areas or naturalized areas on golf courses, or other areas that are not regularly mowed or maintained. Aminopyralid, one of the active ingredients in NativeKlean, has limited activity on woody species when applied to the soil. NativeKlean can be used to treat broadleaf weeds up to the water's edge.

Application precautions around trees

NativeKlean™ herbicide **should NOT** be used over the top of desirable trees. Do not spray NativeKlean™ herbicide within a distance from the base of the tree that is equal to the height of the tree to prevent root uptake.

Do not spray NativeKlean™ herbicide within a distance from the base of the tree that is equal to the height of the tree to prevent root uptake.



Refer to the list of trees below for applications to be made outside of the buffer of the height of the tree:

Common name	Scientific name
Birch	<i>Betula spp.</i>
Black locust	<i>Robinia pseudoacacia</i>
Caragana	<i>Caragana arborescens</i>
Cherry (except black cherry)	<i>Prunus spp.</i>
Chinese elm	<i>Ulmus parviflora</i>
Crabapple	<i>Malus spp.</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
Fir	<i>Abies spp.</i>
Hackberry	<i>Celtis spp.</i>
Honeylocust	<i>Gleditsia triacanthos</i>
Japanese elm	<i>Zelkova serrata</i>
Junipers	<i>Juniperus spp.</i>
Grapes	<i>Vitis spp.</i>
Mimosa	<i>Albizia julibrissin</i>
Mulberry	<i>Morus spp.</i>
Locust species	All species
Lodgepole pine	<i>Pinus contorta</i>
Pinyon pine	<i>Pinus edulis</i>
Ponderosa pine	<i>Pinus ponderosa</i>
Redbud	<i>Cercis occidentalis; or canadensis</i>
Redwoods	<i>Sequoia sempervirens; or related species</i>
Rose	<i>Rosa spp.</i>
Spruce	<i>Picea spp.</i>
Virginia pine	<i>Pinus virginiana</i>
Western red cedar	<i>Thuja spp.</i>

CAUTION: NativeKlean can cause injury or death to desirable vegetation. Users are advised not to apply NativeKlean over the top of desirable trees or in the root zone of susceptible species where injury cannot be tolerated.

Equipment guidelines

Apply NativeKlean™ herbicide with equipment that ensures a coarse low-pressure spray. Equipment used to apply this product should be thoroughly cleaned as follows before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water away from water supplies.
2. Rinse again, this time adding 1 quart of household ammonia or tank-cleaning agent for every 25 gallons of water. Circulate the solution through the entire system to ensure all internal surfaces are contacted for roughly 15 to 20 minutes. Let the solution stand for several hours, preferably overnight.
3. After letting the solution stand for several hours (or overnight), flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Spray nozzles and screens should be removed and cleaned separately.